

REALIZATION OF GHZ STATES AND THE GHZ TEST VIA CAVITY QED FOR A CAVITY PREPARED IN A SUPERPOSITION OF ZERO AND ONE FOCK STATES

E. S. Guerra

Departamento de Física

Universidade Federal Rural do Rio de Janeiro

Cx. Postal 23851, 23890-000 Seropédica, RJ, Brazil

email: emerson@ufrj.br

February 1, 2008

Abstract

In this article we discuss the realization of atomic GHZ states involving three-level atoms in a cascade and in a lambda configuration and we show explicitly how to use this state to perform the GHZ test in which it is possible to decide between local realism theories and quantum mechanics. The experimental realizations proposed makes use the interaction of Rydberg atoms with a cavity prepared in a state which is a superposition of zero and one Fock states.

PACS: 03.65.Ud; 03.67.Mn; 32.80.-t; 42.50.-p

Keywords: GHZ states; cavity QED.

1 INTRODUCTION

The superposition principle, entanglement and its consequence non-locality makes quantum mechanics fascinating and, at the time, intriguing and not intuitive according to our common sense which is based essentially on classical mechanics. Quantum mechanics has given rise to many questions of philosophical nature. For instance, the concept of reality [1, 2, 3, 4, 5] has new interpretations under the light of quantum physics. The Consequences of entanglement and non-locality were noticed by Einstein, Podolsky and Rosen who proposed a *gedanken* experiment, the EPR experiment in a very interesting article [6, 2, 3, 4, 7] to show that quantum theory was not a complete theory. In their article they describe an experimental arrangement involving correlated pair of particles where these particles interact and then are separated. .Therefore, measurements made on one particle can be used via correlation to generate predictions about the other particle. EPR state that "If, without in any way disturbing a system we can predict with certainty the value of a physical quantity, then there exists an element of physical reality corresponding to this physical quantity" and as a necessary condition for the completeness of the theory, EPR state that "every element of the physical reality

must have a counter part in the physical theory". EPR noticed that the fact that quantum mechanics admits no dispersion-free states, does not tell us whether the theory is complete or not. In their words: "From [the dispersion principle] it follows (1) the quantum mechanics description of reality given by the wave function is not complete or (2) when operators corresponding to two physical quantities do not commute the two quantities cannot have simultaneous reality". Shortly after the EPR paper Bohr published a reply in which he defends the completeness of the quantum-mechanical description of nature which could not be refuted properly by EPR and which came to be known as the Copenhagen interpretation [8].

Hidden variable theories [2, 3] were proposed in which one would recover local realism and one of its proponents was Bohm [9, 2, 3]. Bell developed a clever theoretical tool which could be used to perform experiments based on it which could decide between local realism theories and quantum mechanics, the Bell's inequality [10, 2, 3, 7]. In the 1980s it was realized experiments by Aspect and collaborators [11, 7] based on the Bell's inequality which although strongly favored quantum mechanics, there remained some possibility for which a local reality view could still be maintained. We should point out that the main difficulty in the test of the Bell's inequality is related to the fact that there is no perfect particle detectors. For a review of Bell's inequalities and some of its variants see [12].

In 1989, Greenberger, Horne and Zeilinger found out an ingenious theoretical tool to test quantum mechanics confronted with local hidden variable theories, the Bell theorem without inequalities [13, 14, 12], which can demonstrate the spookiness of quantum mechanics even more dramatically than the Bell's analysis. In the GHZ test the decision between quantum mechanics and local realism theories is a binary simple one showing the power of the GHZ formalism. There are several proposals of preparation of GHZ states. For instance, proposals involving cavity QED are presented in [17, 15, 18, 16]. For a concrete experimental realization of GHZ states see [19, 16]. In reference [17] we have proposed a preparation of atomic GHZ states and a GHZ test similar the one discussed in this article. However, in reference [17] we have used three-level lambda atoms in which the lower states are degenerate states interacting with a cavity prepared in a coherent state. In the present article we assume that the atoms are Rydberg atoms of relatively long radiative lifetimes [20] and that the superconducting microwave cavities [21, 22] are perfect cavities, that is, we neglect effects due to decoherence. We are going to consider cavities which are prepared in a superposition of zero and one Fock states.

2 ATOMS IN A CASCADE CONFIGURATION

We start showing how to prepare a cavity C in the state

$$|\psi\rangle_C = \frac{(|0\rangle + |1\rangle)}{\sqrt{2}}. \quad (2.1)$$

In order to prepare this state, we send a two-level atom $A0$, with $|f_0\rangle$ and $|e_0\rangle$ being the lower and upper level respectively, through a Ramsey cavity $R0$ (see (A.72)) in the lower state $|f_0\rangle$ where the atomic states are rotated according to

$$R_0 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & i \\ i & 1 \end{bmatrix}, \quad (2.2)$$

that is,

$$|f_0\rangle \rightarrow \frac{1}{\sqrt{2}}(i|e_0\rangle + |f_0\rangle), \quad (2.3)$$

and through C , resonant with the cavity. Under the Jaynes-Cummings dynamics [23] (see (A.67), for $\Delta = 0$). For $gt = \pi/2$ we see that the state $|f_0\rangle|0\rangle$ does not evolve, however, the state $|e_0\rangle|0\rangle$ evolves to $-i|f_0\rangle|1\rangle$. Then, for the cavity initially in the vacuum state $|0\rangle$, we have

$$\frac{(|f_0\rangle + i|e_0\rangle)}{\sqrt{2}}|0\rangle \longrightarrow |f_0\rangle \frac{(|0\rangle + |1\rangle)}{\sqrt{2}} = |f_0\rangle|\psi\rangle_C. \quad (2.4)$$

Now let us consider a three-level cascade atom Ak with $|e_k\rangle, |f_k\rangle$ and $|g_k\rangle$ being the upper, intermediate and lower atomic state respectively (see Fig. 1). We assume that the transition $|f_k\rangle \rightleftharpoons |e_k\rangle$ is far enough from resonance with the cavity central frequency such that only virtual transitions occur between these states (only these states interact with field in cavity C). In addition we assume that the transition $|e_k\rangle \rightleftharpoons |g_k\rangle$ is highly detuned from the cavity frequency so that there will be no coupling with the cavity field. Here we are going to consider the effect of the atom-field interaction taking into account only levels $|f_k\rangle$ and $|g_k\rangle$. We do not consider level $|e_k\rangle$ since it will not play any role in our scheme. Therefore, we have effectively a two-level system involving states $|f_k\rangle$ and $|g_k\rangle$. Considering levels $|f_k\rangle$ and $|g_k\rangle$, we can write an effective time evolution operator

$$U_k(t) = e^{i\varphi a^\dagger a} |f_k\rangle\langle f_k| + |g_k\rangle\langle g_k|, \quad (2.5)$$

(see (A.70)) where the second term above was put by hand just in order to take into account the effect of level $|g_k\rangle$ and where $\varphi = g^2\tau/\Delta$, g is the coupling constant, $\Delta = \omega_e - \omega_f - \omega$ is the detuning where ω_e and ω_f are the frequencies of the upper and intermediate levels respectively and ω is the cavity field frequency and τ is the atom-field interaction time. Let us take $\varphi = \pi$. Now, let us assume that we let atom $A1$ to interact with cavity C prepared in the state (2.1). Let us assume that atom $A1$ is sent to a Ramsey cavity $R1$ in the lower state $|g_1\rangle$ where it is prepared cprepared in a coherent superposition according to the rotation matrix

$$R_1 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}, \quad (2.6)$$

and we have

$$|\psi\rangle_{A1} = \frac{1}{\sqrt{2}}(|f_1\rangle + |g_1\rangle). \quad (2.7)$$

Taking into account (2.5), after atom $A1$ has passed through the cavity prepared in state (2.1), we get

$$|\psi\rangle_{A1-C} = \frac{1}{2}[(|f_1\rangle + |g_1\rangle)|0\rangle + (-|f_1\rangle + |g_1\rangle)|1\rangle], \quad (2.8)$$

Now, if atom $A1$ enters a second Ramsey cavity $R2$ where the atomic states are rotated according to the rotation matrix

$$R_2 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}, \quad (2.9)$$

we have

$$\begin{aligned} \frac{1}{\sqrt{2}}(|f_1\rangle + |g_1\rangle) &\rightarrow |f_1\rangle, \\ \frac{1}{\sqrt{2}}(-|f_1\rangle + |g_1\rangle) &\rightarrow |g_1\rangle, \end{aligned} \quad (2.10)$$

and, therefore,

$$|\psi\rangle_{A1-C} = \frac{1}{\sqrt{2}}[|f_1\rangle|0\rangle + |g_1\rangle|1\rangle], \quad (2.11)$$

Now, let us prepare a two-level atom $A2$ in the Ramsey cavity $R3$. If atom $A2$ is initially in the state $|g_2\rangle$, according to the rotation matrix

$$R_3 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}, \quad (2.12)$$

we have

$$|\psi\rangle_{A2} = \frac{1}{\sqrt{2}}(|f_2\rangle + |g_2\rangle), \quad (2.13)$$

and let us send this atom through cavity C , assuming that for atom $A2$, as above for atom $A1$, the transition $|f_2\rangle \Rightarrow |e_2\rangle$ is highly detuned from the cavity central frequency. Taking into account (2.5), after the atom has passed through the cavity we get

$$|\psi\rangle_{A1-A2-C} = \frac{1}{2}[|f_1\rangle(|f_2\rangle + |g_2\rangle)|0\rangle + |g_1\rangle(-|f_2\rangle + |g_2\rangle)|1\rangle], \quad (2.14)$$

Then, atom $A2$ enters a Ramsey cavity $R4$ where the atomic states are rotated according to the rotation matrix

$$R_4 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}, \quad (2.15)$$

that is,

$$\begin{aligned} \frac{1}{\sqrt{2}}(|f_2\rangle + |g_2\rangle) &\rightarrow |f_2\rangle, \\ \frac{1}{\sqrt{2}}(-|f_2\rangle + |g_2\rangle) &\rightarrow |g_2\rangle, \end{aligned} \quad (2.16)$$

and we get

$$|\psi, +\rangle_{A1-A2-C} = \frac{1}{\sqrt{2}}[|f_1\rangle |f_2\rangle |0\rangle + |g_1\rangle |g_2\rangle |1\rangle], \quad (2.17)$$

which is a GHZ state. If we had started with the cavity in the state $(|0\rangle - |1\rangle)/\sqrt{2}$ we would get the GHZ state

$$|\psi, -\rangle_{A1-A2-C} = \frac{1}{\sqrt{2}}[|f_1\rangle |f_2\rangle |0\rangle - |g_1\rangle |g_2\rangle |1\rangle]. \quad (2.18)$$

Let us now first discuss a summary of the GHZ test prescription. We will follow closely the discussion presented in the very clear article by Mermin [14]. First we define the atomic operators

$$\begin{aligned} A &= \sigma_x^1 \sigma_y^2 \sigma_y^3, \\ B &= \sigma_y^1 \sigma_x^2 \sigma_y^3, \\ C &= \sigma_y^1 \sigma_y^2 \sigma_x^3, \\ D &= \sigma_x^1 \sigma_x^2 \sigma_x^3, \end{aligned} \quad (2.19)$$

where

$$\sigma_x^k = |f_k\rangle\langle g_k| + |g_k\rangle\langle f_k|, \quad (2.20)$$

$$\sigma_y^k = -i(|f_k\rangle\langle g_k| - |g_k\rangle\langle f_k|), \quad (2.21)$$

$$\sigma_x^3 = |0\rangle\langle 1| + |1\rangle\langle 0|, \quad (2.22)$$

$$\sigma_y^3 = -i(|0\rangle\langle 1| - |1\rangle\langle 0|), \quad (2.22)$$

($k = 1, 2$). It is easy to show that the commutators

$$[A, B] = [A, C] = [B, C] = 0, \quad (2.23)$$

and that

$$A | \psi, \pm \rangle_{A1-A2-C} = B | \psi, \pm \rangle_{A1-A2-C} = C | \psi, \pm \rangle_{A1-A2-C} = \mp 1 | \psi, \pm \rangle_{A1-A2-C}, \quad (2.24)$$

and

$$D | \psi, \pm \rangle_{A1-A2-C} = \pm 1 | \psi, \pm \rangle_{A1-A2-C}. \quad (2.25)$$

If we assume that there are six elements of reality m_x^k and m_y^k ($k = 1, 2$ and 3) each having value $+1$ or -1 waiting to be revealed according to a local realism theory, then we can write

$$\begin{aligned} a_{\pm} &= m_x^1 m_y^2 m_y^3 = \mp 1, \\ b_{\pm} &= m_y^1 m_x^2 m_y^3 = \mp 1, \\ c_{\pm} &= m_y^1 m_y^2 m_x^3 = \mp 1, \end{aligned} \quad (2.26)$$

and we have

$$d_{\pm} = a_{\pm} b_{\pm} c_{\pm} = m_x^1 m_x^2 m_x^3 = \mp 1, \quad (2.27)$$

where the upper and lower subindexes refer to the GHZ state (2.17) and (2.18) respectively and we have used $(m_y^k)^2 = 1$. So, the existence of elements of reality implies that if we measure the value of the observables σ_x^k ($k = 1, 2$ and 3) (that is the elements of reality associated with them) in the state $| \psi, \pm \rangle_{A1-A2-C}$ the product of the three resulting values must be $d_{\pm} = \mp 1$. But according to (2.25) the eigenvalue of the operator D applied the state $| \psi, \pm \rangle_{A1-A2-C}$ is ± 1 . Therefore, measuring this eigenvalue we can decide between theories based on local realism and quantum mechanics.

Now, let us see how we proceed to perform the GHZ test (we follow closely the scheme presented in [17]). We start letting atoms Ak to pass through the Ramsey zones Kk ($k = 1, 2$) where the atomic states are rotated according to the rotation matrix

$$K_k = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}, \quad (2.28)$$

or

$$K_k = \frac{1}{\sqrt{2}} (| f_k \rangle \langle f_k | - | f_k \rangle \langle g_k | + | g_k \rangle \langle f_k | + | g_k \rangle \langle g_k |). \quad (2.29)$$

These rotation matrixes are the key ingredient which allow us to perform the GHZ test. The method of the test we are going to describe is based on a gradual unraveling of the GHZ state being considered.

The eigenvalues of the operators σ_x^k are

$$| \psi_x^k, \pm \rangle = \frac{1}{\sqrt{2}} (| f_k \rangle \pm | g_k \rangle). \quad (2.30)$$

for $k = 1, 2$. Let us take the state (2.17). Writing the states $| f_1 \rangle$ and $| g_1 \rangle$ in terms of the states (2.30) for $k = 1$ and substituting in the GHZ state (2.17) we get

$$| \psi, + \rangle_{A1-A2-C} = \frac{1}{2} [| \psi_x^1, + \rangle (| f_2 \rangle | 0 \rangle + | g_2 \rangle | 1 \rangle) + | \psi_x^1, - \rangle (| f_2 \rangle | 0 \rangle - | g_2 \rangle | 1 \rangle)]. \quad (2.31)$$

Applying (2.29) in the state (2.17) for $k = 1$, we have

$$K_1 |\psi, +\rangle_{A1-A2-C} = \frac{1}{2} [|f_1\rangle(|f_2\rangle|0\rangle - |g_2\rangle|1\rangle) + |g_1\rangle(|f_2\rangle|0\rangle + |g_2\rangle|1\rangle)]. \quad (2.32)$$

Now, we compare (2.32) and (2.31). We see that the rotation by K_1 followed by the detection of $|g_1\rangle$ corresponds to the detection of the state $|\psi_x^1, +\rangle$ whose eigenvalue of σ_x^1 is $+1$. After we detect $|g_1\rangle$, we get

$$|\psi\rangle_{A2-C} = \frac{1}{\sqrt{2}}(|f_2\rangle|0\rangle + |g_2\rangle|1\rangle). \quad (2.33)$$

As before, we rewrite the states $|f_2\rangle$ and $|g_2\rangle$ in terms of these states (2.30) for $k = 2$ and substitute in the above state and we have

$$|\psi\rangle_{A2-C} = \frac{1}{2} [|\psi_x^2, +\rangle(|0\rangle + |1\rangle) + |\psi_x^2, -\rangle(|0\rangle - |1\rangle)]. \quad (2.34)$$

If we apply (2.29) for $k = 2$ to the state (2.33) we get

$$K_2 |\psi\rangle_{A2-C} = \frac{1}{2} [|f_2\rangle(|0\rangle - |1\rangle) + |g_2\rangle(|0\rangle + |1\rangle)]. \quad (2.35)$$

Again, if we compare (2.35) with (2.34), we see that the rotation by K_2 followed by the detection of $|g_2\rangle$ corresponds to the detection of the state $|\psi_x^2, +\rangle$ whose eigenvalue of σ_x^2 is $+1$. After we detect $|g_2\rangle$, we get

$$|\psi_x^3, +\rangle = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle), \quad (2.36)$$

which is the eigenvector of σ_x^3 with eigenvalue $+1$. We now send a two-level atom $A3$, resonant with the cavity, with $|g_3\rangle$ and $|f_3\rangle$ being the lower and upper level respectively, through C . If $A3$ is sent in the lower state $|g_3\rangle$, under the Jaynes-Cummings dynamics, for $gt = \pi/2$, we know that the state $|g_3\rangle|0\rangle$ does not evolve, however, the state $|g_3\rangle|1\rangle$ evolves to $-i|f_3\rangle|0\rangle$. Then we get

$$|g_3\rangle|\psi_x^3, +\rangle \longrightarrow \frac{(|g_3\rangle - i|f_3\rangle)}{\sqrt{2}}|0\rangle \quad (2.37)$$

Now we let atom $A3$ to enter a Ramsey cavity $K3$ where the atomic states are rotated according to the rotation matrix

$$K_3 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & i \\ i & 1 \end{bmatrix}, \quad (2.38)$$

that is,

$$\frac{1}{\sqrt{2}}(-i|f_3\rangle + |g_3\rangle) \rightarrow |g_3\rangle, \quad (2.39)$$

and we see that the rotation by K_3 followed by the detection of $|g_3\rangle$ corresponds to the detection of the state $|\psi_x^2, +\rangle$ whose eigenvalue of σ_x^3 is $+1$.

In the case of

$$|\psi_x^3, -\rangle = \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle) \quad (2.40)$$

which is the eigenvector of σ_x^3 with eigenvalue -1 , we would get

$$|g_3\rangle|\psi_x^3, -\rangle \longrightarrow \frac{(|g_3\rangle + i|f_3\rangle)}{\sqrt{2}}|0\rangle \quad (2.41)$$

and after the rotation (2.38) we would have

$$\frac{1}{\sqrt{2}}(i | f_3 \rangle + | g_3 \rangle) \rightarrow i | f_3 \rangle. \quad (2.42)$$

and we see that the rotation by K_3 followed by the detection of $|f_3\rangle$ corresponds to the detection of the state $|\psi_x^2, -\rangle$ whose eigenvalue of σ_x^3 is -1 .

We can repeat the above procedure and see that we have only four possibilities which are presented schematically below, where on the left, we present the possible sequences of atomic state rotations through K_k and detections of $|f_j\rangle$ or $|g_j\rangle$ ($j = 1, 2$) and $|g_3\rangle$ or $|f_3\rangle$ and on the right, we present the sequences of the corresponding states $|\psi_x^k, \pm\rangle$ where $k = 1, 2$ and 3 which corresponds to the measurement of the eigenvalue of the operator D given by (2.19),

$$\begin{aligned} (K_1, |g_1\rangle)(K_2, |g_2\rangle)(K_3, |g_3\rangle) &\longleftrightarrow |\psi_x^1, +\rangle |\psi_x^2, +\rangle |\psi_x^3, +\rangle, \\ (K_1, |g_1\rangle)(K_2, |f_2\rangle)(K_3, |f_3\rangle) &\longleftrightarrow |\psi_x^1, +\rangle |\psi_x^2, -\rangle |\psi_x^3, -\rangle, \\ (K_1, |f_1\rangle)(K_2, |f_2\rangle)(K_3, |g_3\rangle) &\longleftrightarrow |\psi_x^1, -\rangle |\psi_x^2, -\rangle |\psi_x^3, +\rangle, \\ (K_1, |f_1\rangle)(K_2, |g_2\rangle)(K_3, |f_3\rangle) &\longleftrightarrow |\psi_x^1, -\rangle |\psi_x^2, +\rangle |\psi_x^3, -\rangle. \end{aligned} \quad (2.43)$$

Notice that all these results give us the eigenvalue $+1$ of the operator D (see (2.25)). Therefore, in one run, if we succeed to perform the above rotations of the atomic states according to (2.29), each one followed by the detection of the respective state $|f_j\rangle$ or $|g_j\rangle$ ($j = 1, 2$) and $|g_3\rangle$ or $|f_3\rangle$, we get the result of the experiment in favor of quantum mechanics.

If we perform the test for the state (2.18), following the above procedure, it is easy to see that we have the four possible outcomes,

$$\begin{aligned} (K_1, |g_1\rangle)(K_2, |g_2\rangle)(K_3, |f_3\rangle) &\longleftrightarrow |\psi_x^1, +\rangle |\psi_x^2, +\rangle |\psi_x^3, -\rangle, \\ (K_1, |g_1\rangle)(K_2, |f_2\rangle)(K_3, |g_3\rangle) &\longleftrightarrow |\psi_x^1, +\rangle |\psi_x^2, -\rangle |\psi_x^3, +\rangle, \\ (K_1, |f_1\rangle)(K_2, |f_2\rangle)(K_3, |f_3\rangle) &\longleftrightarrow |\psi_x^1, -\rangle |\psi_x^2, -\rangle |\psi_x^3, -\rangle, \\ (K_1, |f_1\rangle)(K_2, |g_2\rangle)(K_3, |g_3\rangle) &\longleftrightarrow |\psi_x^1, -\rangle |\psi_x^2, +\rangle |\psi_x^3, +\rangle. \end{aligned} \quad (2.44)$$

Notice that all these results give us the eigenvalue -1 of the operator D (see (2.25)), as it should be. See Fig. 2 where we present a scheme of the GHZ experiment.

3 ATOMS IN A LAMBDA CONFIGURATION

Consider a three-level lambda atom (see Fig. 3) interacting with the electromagnetic field inside a cavity C . The states of the atom, $|a\rangle$, $|b\rangle$ and $|c\rangle$ are so that the $|a\rangle \rightleftharpoons |c\rangle$ and $|a\rangle \rightleftharpoons |b\rangle$ transitions are in the far off resonance interaction limit. The time evolution operator for the atom-field interaction $U(t)$ is given by [24]

$$U(\tau) = -e^{i\varphi a^\dagger a} |a\rangle \langle a| + \frac{1}{2}(e^{i\varphi a^\dagger a} + 1) |b\rangle \langle b| + \frac{1}{2}(e^{i\varphi a^\dagger a} - 1) |b\rangle \langle c| + \frac{1}{2}(e^{i\varphi a^\dagger a} - 1) |c\rangle \langle b| + \frac{1}{2}(e^{i\varphi a^\dagger a} + 1) |c\rangle \langle c|, \quad (3.45)$$

where a ($a^\dagger$) is the annihilation (creation) operator for the field in cavity C , $\varphi = 2g^2\tau/\Delta$, g is the coupling constant, $\Delta = \omega_a - \omega_b - \omega = \omega_a - \omega_c - \omega$ is the detuning where ω_a , ω_b and ω_c are the

frequency of the upper level and of the two degenerate lower levels respectively and ω is the cavity field frequency and τ is the atom-field interaction time. For $\varphi = \pi$, we get

$$U(\tau) = -\exp(i\pi a^\dagger a) |a\rangle\langle a| + \Pi_+ |b\rangle\langle b| + \Pi_- |b\rangle\langle c| + \Pi_- |c\rangle\langle b| + \Pi_+ |c\rangle\langle c|, \quad (3.46)$$

where

$$\begin{aligned} \Pi_+ &= \frac{1}{2}(e^{i\pi a^\dagger a} + 1), \\ \Pi_- &= \frac{1}{2}(e^{i\pi a^\dagger a} - 1). \end{aligned} \quad (3.47)$$

Let us first show how we can get a GHZ state making use of three-level lambda atoms interacting with a cavity field prepared in state (2.1). Consider the atom $A1$ in the state $|\psi\rangle_{A1} = |b_1\rangle$ and a cavity C prepared in the state (2.1). We now let atom $A1$ fly through the cavity C . Taking into account (3.46) the state of the system $A1 - C$ evolves to

$$|\psi\rangle_{A1-C} = \frac{1}{\sqrt{2}}(|b_1\rangle|0\rangle - |c_1\rangle|1\rangle). \quad (3.48)$$

Consider now another three-level lambda atom $A2$ prepared initially in the state $|b_2\rangle$, which are going to pass through the cavity. After this second atom has passed through the cavity, the system evolves to

$$|\psi, +\rangle_{A1-A2-C} = \frac{1}{\sqrt{2}}(|b_1\rangle|b_2\rangle|0\rangle + |c_1\rangle|c_2\rangle|1\rangle). \quad (3.49)$$

which is a a GHZ state. If we had started with the cavity in the state $(|0\rangle - |1\rangle)/\sqrt{2}$ we would get the GHZ state

$$|\psi, -\rangle_{A1-A2-C} = \frac{1}{\sqrt{2}}[|b_1\rangle|b_2\rangle|0\rangle - |c_1\rangle|c_2\rangle|1\rangle]. \quad (3.50)$$

In the case of (3.49), following the prescription of the previous section, we can see that in the GHZ test we have only four possibilities which are presented schematically below, where on the left, we present the possible sequences of atomic state rotations through K_k and detections of $|b_j\rangle$ or $|c_j\rangle$ ($j = 1, 2$) and $|g_3\rangle$ or $|f_3\rangle$ and on the right, we present the sequences of the corresponding states $|\psi_x^k, \pm\rangle$ where $k = 1, 2$ and 3 which corresponds to the measurement of the eigenvalue of the operator D given by (2.19),

$$\begin{aligned} (K_1, |c_1\rangle)(K_2, |c_2\rangle)(K_3, |g_3\rangle) &\longleftrightarrow |\psi_x^1, +\rangle|\psi_x^2, +\rangle|\psi_x^3, +\rangle, \\ (K_1, |c_1\rangle)(K_2, |b_2\rangle)(K_3, |f_3\rangle) &\longleftrightarrow |\psi_x^1, +\rangle|\psi_x^2, -\rangle|\psi_x^3, -\rangle, \\ (K_1, |b_1\rangle)(K_2, |b_2\rangle)(K_3, |g_3\rangle) &\longleftrightarrow |\psi_x^1, -\rangle|\psi_x^2, -\rangle|\psi_x^3, +\rangle, \\ (K_1, |b_1\rangle)(K_2, |c_2\rangle)(K_3, |f_3\rangle) &\longleftrightarrow |\psi_x^1, -\rangle|\psi_x^2, +\rangle|\psi_x^3, -\rangle. \end{aligned} \quad (3.51)$$

where now,

$$|\psi_x^k, \pm\rangle = \frac{1}{\sqrt{2}}(|b_k\rangle \pm |c_k\rangle), \quad (3.52)$$

for $k = 1, 2$. Notice that all these results give us the eigenvalue $+1$ of the operator D (see (2.25)). Therefore, in one run, if we succeed to perform the above rotations of the atomic states according to (2.29), each one followed by the detection of the respective state $|b_j\rangle$ or $|c_j\rangle$ ($j = 1, 2$) and $|g_3\rangle$ or $|f_3\rangle$, we get the result of the experiment in favor of quantum mechanics.

If we perform the test for the state (3.50), following the above procedure, it is easy to see that we have the four possible outcomes,

$$\begin{aligned}
(K_1, |c_1\rangle)(K_2, |c_2\rangle)(K_3, |f_3\rangle) &\longleftrightarrow |\psi_x^1, +\rangle |\psi_x^2, +\rangle |\psi_x^3, -\rangle, \\
(K_1, |c_1\rangle)(K_2, |b_2\rangle)(K_3, |g_3\rangle) &\longleftrightarrow |\psi_x^1, +\rangle |\psi_x^2, -\rangle |\psi_x^3, +\rangle, \\
(K_1, |b_1\rangle)(K_2, |b_2\rangle)(K_3, |f_3\rangle) &\longleftrightarrow |\psi_x^1, -\rangle |\psi_x^2, -\rangle |\psi_x^3, -\rangle, \\
(K_1, |b_1\rangle)(K_2, |c_2\rangle)(K_3, |g_3\rangle) &\longleftrightarrow |\psi_x^1, -\rangle |\psi_x^2, +\rangle |\psi_x^3, +\rangle.
\end{aligned} \tag{3.53}$$

Notice that all these results give us the eigenvalue -1 of the operator D (see (2.25)), as it should be. See Fig. 4 where we present a scheme of the GHZ experiment.

4 CONCLUSION

We have presented two schemes of preparation of GHZ states and realization of the GHZ test via cavity QED. In these schemes the cavity is prepared in a superposition of zero and one Fock states which are not affected dramatically by decoherence and which are relatively easy to handle. In the first scheme we make use of atoms in a cascade configuration and in the second scheme we make use of atoms in a three-level lambda configuration. The advantage of using a cascade atomic configuration is that the detection process is easier, however, we have to perform many atomic state rotations. The advantage of using the lambda atomic configuration is that we do not have to use many atomic state rotations, however, the detection of the atomic states are more complicated.

A Time evolution operator

A.1 Two-level atoms

Let us consider a two-level atom interacting with a cavity field, where $|e\rangle$ and $|f\rangle$ are the upper and lower states respectively, with ω_e and ω_f being the two atomic frequencies associated to these two states and ω the cavity field frequency (see Fig. 1). The Jaynes-Cummings Hamiltonian, under the rotating-wave approximation, is given by

$$H = \hbar a^\dagger a + \hbar \omega_e |e\rangle \langle e| + \hbar \omega_f |f\rangle \langle f| + \hbar g [a|e\rangle \langle f| + a^\dagger |f\rangle \langle e|], \tag{A.54}$$

where $a^\dagger$ and a are the creation and annihilation operators respectively for the cavity field, g is the coupling constant and we write

$$H = H_0 + H_I, \tag{A.55}$$

where we have settled

$$\begin{aligned}
H_0 &= \hbar a^\dagger a + \hbar \omega_e |e\rangle \langle e| + \hbar \omega_f |f\rangle \langle f|, \\
H_I &= \hbar g (a|e\rangle \langle f| + a^\dagger |f\rangle \langle e|).
\end{aligned} \tag{A.56}$$

Lets define the interaction picture

$$|\psi_I\rangle = e^{i\frac{H_0}{\hbar}t} |\psi_S\rangle, \tag{A.57}$$

where

$$i\hbar \frac{d}{dt} |\psi_S\rangle = H |\psi_S\rangle. \quad (\text{A.58})$$

Then, we get

$$i\hbar \frac{d}{dt} |\psi_I\rangle = V_I |\psi_I\rangle, \quad (\text{A.59})$$

where

$$V_I = e^{i\frac{H_0}{\hbar}t} H_I e^{-i\frac{H_0}{\hbar}t} = \hbar \begin{bmatrix} 0 & ge^{i\Delta t}a \\ ge^{-i\Delta t}a^\dagger & 0 \end{bmatrix} \quad (\text{A.60})$$

and

$$\Delta = (\omega_e - \omega_f) - \omega. \quad (\text{A.61})$$

Considering

$$|\psi_I(t)\rangle = U_I(t) |\psi_I(0)\rangle = U_I(t) |\psi_S(0)\rangle, \quad (\text{A.62})$$

we have to solve the Schrödinger's equation for the time evolution operator

$$i\hbar \frac{dU_I}{dt} = V_I U_I, \quad (\text{A.63})$$

where

$$U_I(t) = \begin{bmatrix} u_{ee}(t) & u_{ef}(t) \\ u_{fe}(t) & u_{ff}(t) \end{bmatrix} \quad (\text{A.64})$$

and

$$U_I(0) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}. \quad (\text{A.65})$$

That is,

$$\begin{aligned} i\frac{d}{dt}u_{ee}(t) &= ge^{i\Delta t}au_{ef}(t), \\ i\frac{d}{dt}u_{ef}(t) &= ge^{i\Delta t}au_{ff}(t), \\ i\frac{d}{dt}u_{fe}(t) &= ge^{-i\Delta t}a^\dagger u_{ee}(t), \\ i\frac{d}{dt}u_{ff}(t) &= ge^{-i\Delta t}a^\dagger u_{ef}(t), \end{aligned} \quad (\text{A.66})$$

which can be solved easily using, for instance, Laplace transformation, and we get

$$U_I(t) = \begin{bmatrix} e^{i\frac{\Delta}{2}t}(\cos \mu t - i\frac{\Delta}{2\mu} \sin \mu t) & -ige^{i\frac{\Delta}{2}t}\frac{1}{\mu}(\sin \mu t)a \\ -iga^\dagger e^{-i\frac{\Delta}{2}t}\frac{1}{\mu}(\sin \mu t) & e^{-i\frac{\Delta}{2}t}(\cos \nu t + i\frac{\Delta}{2\nu} \sin \nu t) \end{bmatrix}, \quad (\text{A.67})$$

where we have defined

$$\begin{aligned} \mu &= \sqrt{g^2aa^\dagger + \frac{\Delta^2}{4}}, \\ \nu &= \sqrt{g^2a^\dagger a + \frac{\Delta^2}{4}}. \end{aligned} \quad (\text{A.68})$$

In the large detuning limit ($\Delta \gg g$) we have

$$\begin{aligned}\mu &= \sqrt{g^2 aa^\dagger + \frac{\Delta^2}{4}} \cong \frac{\Delta}{2} + \frac{g^2 aa^\dagger}{\Delta}, \\ \nu &= \sqrt{g^2 a^\dagger a + \frac{\Delta^2}{4}} \cong \frac{\Delta}{2} + \frac{g^2 a^\dagger a}{\Delta}.\end{aligned}\tag{A.69}$$

and we get easily

$$U_d(t) = e^{-i\varphi(a^\dagger a + 1)} |e\rangle\langle e| + e^{i\varphi a^\dagger a} |f\rangle\langle f|,\tag{A.70}$$

where $\varphi = g^2 t / \Delta$. If we neglect spontaneous emission, the above time evolution operator can also be obtained from the effective Hamiltonian

$$H_d = \hbar \frac{g^2}{\Delta} a^\dagger a (|e\rangle\langle e| - |f\rangle\langle f|).\tag{A.71}$$

The subindexes in (A.70) and (A.71) are related to the atom-field interaction described by them, that is, a dispersive interaction.

In the case we have a resonant interaction of an atom with cavity field ($\Delta = 0$ in (A.67)), if the field is a very intense field we can treat it classically. That is, in the time evolution operator (A.67) we set $\Delta = 0$, and we substitute the creation and annihilation field operators according to $a \rightarrow \eta e^{i\theta}$ and $a^\dagger \rightarrow \eta e^{-i\theta}$ where η and θ are c-numbers. Then, we have a semiclassical approach of the atom-field interaction in which the field is treated classically and the atoms according to quantum mechanics. In this case (A.67) becomes

$$U_{I,SC}(t) = \begin{bmatrix} \cos(g\eta t) & -ie^{i\theta} \sin(g\eta t) \\ -ie^{-i\theta} \sin(g\eta t) & \cos(g\eta t) \end{bmatrix}.\tag{A.72}$$

Now we take $\theta = \pi/2$. If we choose $g\eta t = \pi/4$ we have the rotation matrix

$$R_{\frac{\pi}{2}, \frac{\pi}{4}} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix},\tag{A.73}$$

and for $g\eta t = -\pi/4$ we have

$$R_{\frac{\pi}{2}, -\frac{\pi}{4}} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}.\tag{A.74}$$

For $g\eta t = \pi/2$ we get

$$R_{\frac{\pi}{2}, \frac{\pi}{2}} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix},\tag{A.75}$$

and for $g\eta t = -\pi/2$

$$R_{\frac{\pi}{2}, -\frac{\pi}{2}} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}.\tag{A.76}$$

Now if we take $\theta = \pi$ and $g\eta t = \pi/4$ we have the rotation matrix

$$R_{\pi, \frac{\pi}{4}} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & i \\ i & 1 \end{bmatrix},\tag{A.77}$$

Choosing the proper values of θ , g , η and t we can get the rotation matrix we need to perform the rotation of the atomic states we desire in a Ramsey cavity. In all the sections of this article when we mention a rotation of atomic states in a Ramsey cavity we are taking into account (A.72).

Just as a remark, consider (??) and assume we have an intense field, that is, we can use a semiclassical approach. In this case we set $a \rightarrow \eta e^{i\theta}$ and $a^\dagger \rightarrow \eta e^{-i\theta}$ where η and θ are c-numbers and $a^\dagger a \rightarrow \eta^2$. Defining $\varphi a^\dagger a \rightarrow \varphi \eta^2 = \beta/2$, (??) reads

$$U_{d,SC} = \begin{bmatrix} e^{-i\beta/2} & 0 \\ 0 & e^{i\beta/2} \end{bmatrix}, \quad (\text{A.78})$$

and for $\beta = \pi$ we have

$$U_\pi = i \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}. \quad (\text{A.79})$$

Any arbitrary 2×2 unitary matrix M may be decomposed as [?]

$$M = e^{i\alpha} \begin{bmatrix} e^{-i\beta/2} & 0 \\ 0 & e^{i\beta/2} \end{bmatrix} \begin{bmatrix} \cos \frac{\gamma}{2} & -\sin \frac{\gamma}{2} \\ \sin \frac{\gamma}{2} & \cos \frac{\gamma}{2} \end{bmatrix} \begin{bmatrix} e^{-i\delta/2} & 0 \\ 0 & e^{i\delta/2} \end{bmatrix}, \quad (\text{A.80})$$

where α, β, γ and δ are real parameters. Therefore, we can use (A.72) and (A.78) to get a rotation matrix we need.

A.2 Three-level lambda atoms

We start with the Hamiltonian of a degenerate three-level lambda atom (see Fig. 3) interacting with a field cavity mode

$$H = \hbar\omega a^\dagger a + \hbar\omega_a |a\rangle\langle a| + \hbar\omega_b |b\rangle\langle b| + \hbar\omega_c |c\rangle\langle c| + \hbar a(g_1 |a\rangle\langle b| + g_2 |a\rangle\langle c|) + \hbar a^\dagger(g_1^* |b\rangle\langle a| + g_2^* |c\rangle\langle a|), \quad (\text{A.81})$$

where $|a\rangle$, $|b\rangle$ and $|c\rangle$ are the upper and the two degenerated lower atomic levels respectively, a ($a^\dagger$) is the annihilation (creation) field operator and g_1 and g_2 are the coupling constants corresponding to the transitions $|a\rangle \rightleftharpoons |c\rangle$ and $|a\rangle \rightleftharpoons |b\rangle$, respectively. In the interaction picture,

$$V = \hbar \left[e^{i\Delta_1 t} g_1 a |a\rangle\langle b| + e^{-i\Delta_1 t} g_1^* a^\dagger |b\rangle\langle a| \right] + \hbar \left[e^{i\Delta_2 t} g_2 a |a\rangle\langle c| + e^{-i\Delta_2 t} g_2^* a^\dagger |c\rangle\langle a| \right], \quad (\text{A.82})$$

where

$$\begin{aligned} \Delta_1 &= \omega_a - \omega_b - \omega \\ \Delta_2 &= \omega_a - \omega_c - \omega. \end{aligned} \quad (\text{A.83})$$

The time evolution operator

$$U(t) = \begin{bmatrix} u_{aa} & u_{ab} & u_{ac} \\ u_{ba} & u_{bb} & u_{bc} \\ u_{ca} & u_{cb} & u_{cc} \end{bmatrix}, \quad (\text{A.84})$$

whose initial condition is given by

$$U(0) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad (\text{A.85})$$

should satisfy the Schrödinger equation of motion

$$i\hbar \frac{dU}{dt} = VU = \begin{bmatrix} \zeta_1 u_{ba} + \zeta_2 u_{ca} & \zeta_1 u_{bb} + \zeta_2 u_{cb} & \zeta_1 u_{bc} + \zeta_2 u_{cc} \\ \zeta_1^\dagger u_{aa} & \zeta_1^\dagger u_{ab} & \zeta_1^\dagger u_{ac} \\ \zeta_2^\dagger u_{aa} & \zeta_2^\dagger u_{ab} & \zeta_2^\dagger u_{ac} \end{bmatrix}, \quad (\text{A.86})$$

where

$$\begin{aligned}\zeta_1 &= \hbar e^{i\Delta_1 t} g_1 a, \\ \zeta_2 &= \hbar e^{i\Delta_2 t} g_2 a.\end{aligned}\tag{A.87}$$

Observe that this equation may be grouped in three sets of couple differential equations. One for u_{aa} , u_{ba} and u_{ca} , i.e.,

$$\begin{aligned}i\hbar \frac{du_{aa}}{dt} &= \zeta_1 u_{ba} + \zeta_2 u_{ca}, \\ i\hbar \frac{du_{ba}}{dt} &= \zeta_1^\dagger u_{aa}, \\ i\hbar \frac{du_{ca}}{dt} &= \zeta_2^\dagger u_{aa},\end{aligned}\tag{A.88}$$

another involving only u_{ab} , u_{bb} and u_{cb} ,

$$\begin{aligned}i\hbar \frac{du_{ab}}{dt} &= \zeta_1 u_{bb} + \zeta_2 u_{cb}, \\ i\hbar \frac{du_{bb}}{dt} &= \zeta_1^\dagger u_{ab}, \\ i\hbar \frac{du_{cb}}{dt} &= \zeta_2^\dagger u_{ab},\end{aligned}\tag{A.89}$$

and, finally, one involving u_{ac} , u_{bc} and u_{cc} ,

$$\begin{aligned}i\hbar \frac{du_{ac}}{dt} &= \zeta_1 u_{bc} + \zeta_2 u_{cc}, \\ i\hbar \frac{du_{bc}}{dt} &= \zeta_1^\dagger u_{ac}, \\ i\hbar \frac{du_{cc}}{dt} &= \zeta_2^\dagger u_{ac}.\end{aligned}\tag{A.90}$$

Let us take $\Delta_1 = \Delta_2 = \Delta = -i\eta$

$$\begin{aligned}\zeta_1 &= \hbar e^{i\Delta_1 t} g_1 a = \hbar \alpha_1 e^{\eta t}, \\ \zeta_2 &= \hbar e^{i\Delta_2 t} g_2 a = \hbar \alpha_2 e^{\eta t}.\end{aligned}\tag{A.91}$$

Notice that all the 3 system of differential equations above are of the form

$$\begin{aligned}i \frac{dx}{dt} &= \alpha_1 e^{\eta t} y + \alpha_2 e^{\eta t} z, \\ i e^{\eta t} \frac{dy}{dt} &= \alpha_1^\dagger x, \\ i e^{\eta t} \frac{dz}{dt} &= \alpha_2^\dagger x.\end{aligned}\tag{A.92}$$

We can solve the systems of differential equations using, for instance, Laplace transformation and we have for the degenerate case

$$u_{aa}(t) = \frac{e^{i\frac{\Delta t}{2}}}{\sqrt{\mu}} \left[\sqrt{\mu} \cos \sqrt{\mu} t - i \frac{\Delta}{2} \sin \sqrt{\mu} t \right],$$

$$\begin{aligned}
u_{ab}(t) &= -i \frac{e^{i\frac{\Delta t}{2}}}{\sqrt{\mu}} \sin \sqrt{\mu} t \alpha_1, \\
u_{ac}(t) &= -i \frac{e^{i\frac{\Delta t}{2}}}{\sqrt{\mu}} \sin \sqrt{\mu} t \alpha_2, \\
u_{ba}(t) &= -i \alpha_1^\dagger \frac{e^{-i\frac{\Delta t}{2}}}{\sqrt{\mu}} \sin \sqrt{\mu} t, \\
u_{bb}(t) &= 1 + \alpha_1^\dagger \alpha_1 \frac{1}{\sqrt{\nu}} \frac{1}{\alpha_1^\dagger \alpha_1 + \alpha_2^\dagger \alpha_2} \left[e^{-i\frac{\Delta}{2}t} \left(i \frac{\Delta}{2} \sin \sqrt{\nu} t + \sqrt{\nu} \cos \sqrt{\nu} t \right) - \sqrt{\nu} \right], \\
u_{bc}(t) &= \alpha_1^\dagger \alpha_2 \frac{1}{\alpha_1^\dagger \alpha_1 + \alpha_2^\dagger \alpha_2} \left[e^{-i\frac{\Delta}{2}t} \left(i \frac{\Delta}{2\sqrt{\nu}} \sin \sqrt{\nu} t + \cos \sqrt{\nu} t \right) - 1 \right], \\
u_{ca}(t) &= -i \alpha_2^\dagger \frac{e^{-i\frac{\Delta t}{2}}}{\sqrt{\mu}} \sin \sqrt{\mu} t, \\
u_{cb}(t) &= \alpha_2^\dagger \alpha_1 \frac{1}{\alpha_1^\dagger \alpha_1 + \alpha_2^\dagger \alpha_2} \left[e^{-i\frac{\Delta}{2}t} \left(i \frac{\Delta}{2\sqrt{\nu}} \sin \sqrt{\nu} t + \cos \sqrt{\nu} t \right) - 1 \right], \\
u_{cc}(t) &= 1 + \alpha_2^\dagger \alpha_2 \frac{1}{\sqrt{\nu}} \frac{1}{\alpha_1^\dagger \alpha_1 + \alpha_2^\dagger \alpha_2} \left[e^{-i\frac{\Delta}{2}t} \left(i \frac{\Delta}{2} \sin \sqrt{\nu} t + \sqrt{\nu} \cos \sqrt{\nu} t \right) - \sqrt{\nu} \right], \quad (\text{A.93})
\end{aligned}$$

where

$$\begin{aligned}
\mu &= \frac{\Delta^2}{4} + \alpha_1 \alpha_1^\dagger + \alpha_2 \alpha_2^\dagger, \\
\nu &= \frac{\Delta^2}{4} + \alpha_1^\dagger \alpha_1 + \alpha_2^\dagger \alpha_2. \quad (\text{A.94})
\end{aligned}$$

It is easy to show that for the non-degenerate case, i.e.,

$$\begin{aligned}
\alpha_1 &= g_1 a_1, \\
\alpha_2 &= g_2 a_2, \quad (\text{A.95})
\end{aligned}$$

we obtain

$$\begin{aligned}
u_{aa}(t) &= \frac{e^{i\frac{\Delta t}{2}}}{\sqrt{\mu}} \left[\sqrt{\mu} \cos \sqrt{\mu} t - i \frac{\Delta}{2} \sin \sqrt{\mu} t \right], \\
u_{ab}(t) &= -i \frac{e^{i\frac{\Delta t}{2}}}{\sqrt{\mu}} \sin \sqrt{\mu} t \alpha_1, \\
u_{ac}(t) &= -i \frac{e^{i\frac{\Delta t}{2}}}{\sqrt{\mu}} \sin \sqrt{\mu} t \alpha_2, \\
u_{ba}(t) &= -i \alpha_1^\dagger \frac{e^{-i\frac{\Delta t}{2}}}{\sqrt{\mu}} \sin \sqrt{\mu} t, \\
u_{bb}(t) &= 1 + \alpha_1^\dagger \alpha_1 \frac{1}{\sqrt{\nu_1}} \frac{1}{\alpha_1^\dagger \alpha_1 + \alpha_2^\dagger \alpha_2} \left[e^{-i\frac{\Delta}{2}t} \left(i \frac{\Delta}{2} \sin \sqrt{\nu_1} t + \sqrt{\nu_1} \cos \sqrt{\nu_1} t \right) - \sqrt{\nu_1} \right], \\
u_{bc}(t) &= \alpha_1^\dagger \alpha_2 \frac{1}{\alpha_1^\dagger \alpha_1 + \alpha_2^\dagger \alpha_2} \left[e^{-i\frac{\Delta}{2}t} \left(i \frac{\Delta}{2\sqrt{\nu_1}} \sin \sqrt{\nu_1} t + \cos \sqrt{\nu_1} t \right) - 1 \right], \\
u_{ca}(t) &= -i \alpha_2^\dagger \frac{e^{-i\frac{\Delta t}{2}}}{\sqrt{\mu}} \sin \sqrt{\mu} t,
\end{aligned}$$

$$\begin{aligned}
u_{cb}(t) &= \alpha_2^\dagger \alpha_1 \frac{1}{\alpha_1^\dagger \alpha_1 + \alpha_2 \alpha_2^\dagger} \left[e^{-i\frac{\Delta}{2}t} \left(i \frac{\Delta}{2\sqrt{\nu_1}} \sin \sqrt{\nu_1}t + \cos \sqrt{\nu_1}t \right) - 1 \right], \\
u_{cc}(t) &= 1 + \alpha_2^\dagger \alpha_2 \frac{1}{\sqrt{\nu_2} \alpha_1 \alpha_1^\dagger + \alpha_2^\dagger \alpha_2} \left[e^{-i\frac{\Delta}{2}t} \left(i \frac{\Delta}{2} \sin \sqrt{\nu_2}t + \sqrt{\nu_2} \cos \sqrt{\nu_2}t \right) - \sqrt{\nu_2} \right], \quad (\text{A.96})
\end{aligned}$$

where

$$\begin{aligned}
\mu &= \frac{\Delta^2}{4} + \alpha_1 \alpha_1^\dagger + \alpha_2 \alpha_2^\dagger, \\
\nu_1 &= \frac{\Delta^2}{4} + \alpha_1^\dagger \alpha_1 + \alpha_2 \alpha_2^\dagger, \\
\nu_2 &= \frac{\Delta^2}{4} + \alpha_1 \alpha_1^\dagger + \alpha_2^\dagger \alpha_2. \quad (\text{A.97})
\end{aligned}$$

Returning to the degenerate case, in the large detuning limit, we have

$$\begin{aligned}
\sqrt{\mu} &\approx \frac{\Delta^2}{2} + \frac{\alpha_1 \alpha_1^\dagger + \alpha_2 \alpha_2^\dagger}{\Delta} = \frac{\Delta}{2} + \frac{|g_1|^2 + |g_2|^2}{\Delta} aa^\dagger, \\
\sqrt{\nu} &\approx \frac{\Delta^2}{2} + \frac{\alpha_1^\dagger \alpha_1 + \alpha_2^\dagger \alpha_2}{\Delta} = \frac{\Delta}{2} + \frac{|g_1|^2 + |g_2|^2}{\Delta} a^\dagger a, \quad (\text{A.98})
\end{aligned}$$

and we get

$$\begin{aligned}
u_{aa}(t) &= \exp \left(i \frac{|g_1|^2 + |g_2|^2}{\Delta} t aa^\dagger \right), \\
u_{ab}(t) &= 0, \\
u_{ac}(t) &= 0, \\
u_{ba}(t) &= 0, \\
u_{bb}(t) &= 1 + \frac{|g_1|^2}{|g_1|^2 + |g_2|^2} \left[\exp \left(i \frac{|g_1|^2 + |g_2|^2}{\Delta} t a^\dagger a \right) - 1 \right], \\
u_{bc}(t) &= \frac{g_1^* g_2}{|g_1|^2 + |g_2|^2} \left[\exp \left(i \frac{|g_1|^2 + |g_2|^2}{\Delta} t a^\dagger a \right) - 1 \right], \\
u_{ca}(t) &= 0, \\
u_{cb}(t) &= \frac{g_2^* g_1}{|g_1|^2 + |g_2|^2} \left[\exp \left(i \frac{|g_1|^2 + |g_2|^2}{\Delta} t a^\dagger a \right) - 1 \right], \\
u_{cc}(t) &= 1 + \frac{|g_2|^2}{|g_1|^2 + |g_2|^2} \left[\exp \left(i \frac{|g_1|^2 + |g_2|^2}{\Delta} t a^\dagger a \right) - 1 \right]. \quad (\text{A.99})
\end{aligned}$$

If

$$\begin{aligned}
g_1 &= g e^{i\varphi_1}, \\
g_2 &= g e^{i\varphi_2}, \quad (\text{A.100})
\end{aligned}$$

we finally have

$$u_{aa}(t) = \exp \left(i \frac{2g^2 t}{\Delta} \right) \exp \left(i \frac{2g^2 t}{\Delta} a^\dagger a \right),$$

$$\begin{aligned}
u_{ab}(t) &= 0, \\
u_{ac}(t) &= 0, \\
u_{ba}(t) &= 0, \\
u_{bb}(t) &= \frac{1}{2} \left[\exp \left(i \frac{2g^2 t}{\Delta} a^\dagger a \right) + 1 \right], \\
u_{bc}(t) &= \frac{e^{i(\varphi_1 - \varphi_2)}}{2} \left[\exp \left(i \frac{2g^2 t}{\Delta} a^\dagger a \right) - 1 \right], \\
u_{ca}(t) &= 0, \\
u_{cb}(t) &= \frac{e^{-i(\varphi_1 - \varphi_2)}}{2} \left[\exp \left(i \frac{2g^2 t}{\Delta} a^\dagger a \right) - 1 \right], \\
u_{cc}(t) &= \frac{1}{2} \left[\exp \left(i \frac{2g^2 t}{\Delta} a^\dagger a \right) + 1 \right],
\end{aligned} \tag{A.101}$$

which agrees with the result obtained in [24].

Now, let us considering the classical limit in the three level lambda atom interacting with two dephased modes with the same frequency (see (A.96)). We see that the matrix elements of the evolution operator read as:

$$\begin{aligned}
u_{bb} &= 1 + \epsilon_1^* \epsilon_1 \frac{1}{2 |\epsilon|^2} [e^{-i\Delta t/2} (\cos \sqrt{2 |\epsilon|^2 + \Delta^2/4} t \\
&\quad + i \frac{\Delta}{2 \sqrt{2 |\epsilon|^2 + \Delta^2/4}} \sin \sqrt{2 |\epsilon|^2 + \Delta^2/4} t) - 1], \\
u_{cc} &= 1 + \epsilon_2^* \epsilon_2 \frac{1}{2 |\epsilon|^2} [e^{-i\Delta t/2} (\cos \sqrt{2 |\epsilon|^2 + \Delta^2/4} t \\
&\quad + i \frac{\Delta}{2 \sqrt{2 |\epsilon|^2 + \Delta^2/4}} \sin \sqrt{2 |\epsilon|^2 + \Delta^2/4} t) - 1], \\
u_{cb} &= \epsilon_1^* \epsilon_2 \frac{1}{2 |\epsilon|^2} [e^{-i\Delta t/2} (\cos \sqrt{2 |\epsilon|^2 + \Delta^2/4} t \\
&\quad + i \frac{\Delta}{2 \sqrt{2 |\epsilon|^2 + \Delta^2/4}} \sin \sqrt{2 |\epsilon|^2 + \Delta^2/4} t) - 1], \\
u_{bc} &= \epsilon_1 \epsilon_2^* \frac{1}{2 |\epsilon|^2} [e^{-i\Delta t/2} (\cos \sqrt{2 |\epsilon|^2 + \Delta^2/4} t \\
&\quad + i \frac{\Delta}{2 \sqrt{2 |\epsilon|^2 + \Delta^2/4}} \sin \sqrt{2 |\epsilon|^2 + \Delta^2/4} t) - 1],
\end{aligned} \tag{A.102}$$

where we have used

$$\begin{aligned}
g_1 a_1 &\rightarrow \epsilon_1 = \epsilon e^{i\theta_1}, \\
g_2 a_2 &\rightarrow \epsilon_2 = \epsilon e^{i\theta_2}.
\end{aligned} \tag{A.103}$$

In the high detuning limit we obtain the expression

$$u_{bb} = \frac{1}{2} (e^{i\varphi} + 1),$$

$$\begin{aligned}
u_{cc} &= \frac{1}{2}(e^{i\varphi} + 1), \\
u_{cb} &= \frac{1}{2}e^{-i\phi}(e^{i\varphi} - 1), \\
u_{bc} &= \frac{1}{2}e^{i\phi}(e^{i\varphi} - 1),
\end{aligned} \tag{A.104}$$

where $\phi = \theta_1 - \theta_2$ and $\varphi = 2|\epsilon|^2 t/\Delta$. Choosing $\varphi = \pi/2$ and $\phi = \pi/2$ we get

$$U = \frac{1}{\sqrt{2}}e^{i\pi/4} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}. \tag{A.105}$$

Choosing $\varphi = \pi$ and $\phi = \pi/2$ we get

$$U = i \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}. \tag{A.106}$$

Choosing $\varphi = \pi$ and $\phi = -\pi/2$ we get

$$U = i \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \tag{A.107}$$

Choosing $\varphi = \pi$ and $\phi = \pi$ we get

$$U = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \tag{A.108}$$

Figure Captions

Fig. 1- Energy states scheme of a three-level atom where $|e\rangle$ is the upper state with atomic frequency ω_e , $|f\rangle$ is the intermediate state with atomic frequency ω_f , $|g\rangle$ is the lower state with atomic frequency ω_g and ω is the cavity field frequency and $\Delta = (\omega_e - \omega_f) - \omega$ is the detuning. The transition $|f\rangle \rightleftharpoons |e\rangle$ is far enough of resonance with the cavity central frequency such that only virtual transitions occur between these levels (only these states interact with field in cavity C). In addition we assume that the transition $|e\rangle \rightleftharpoons |g\rangle$ is highly detuned from the cavity frequency so that there will be no coupling with the cavity field in C .

Fig. 2- Set-up for the GHZ experiment. Atom $A0$ is sent to the cavity $R0$, where it is prepared in a coherent superposition, and to cavity C to prepare it in a state which is a superposition of zero and one Fock states. Atom $A1$ passes through the Ramsey cavity $R1$ where it is prepared in a coherent superposition, cavity C and through the Ramsey cavity $R2$. Atom $A2$ passes through the Ramsey cavity $R3$ where it is prepared in a coherent superposition, cavity C and through the Ramsey cavity $R4$. Atom $A3$ is prepared in state $|g_3\rangle$, and sent to the cavity C . Once the GHZ state has been obtained, the GHZ test is performed making use of the Ramsey cavities $K1, K2$ and $K3$ and detectors $D1, D2$ and $D3$ as described in the text.

Fig. 3- Energy level scheme of the three-level lambda atom where $|a\rangle$ is the upper state with atomic frequency ω_a , $|b\rangle$ and $|c\rangle$ are the lower states with atomic frequency ω_b and ω_c , ω is the cavity field frequency and $\Delta = \omega_a - \omega_b - \omega = \omega_a - \omega_c - \omega$ is the detuning.

Fig. 4- Set-up for the GHZ experiment. Atom $A0$ is sent to the cavity $R0$, where it is prepared in a coherent superposition, and to cavity C to prepare it in a state which is a superposition of zero and one Fock states. Atom $A1$ is prepared in state $|b_1\rangle$ and passes through cavity C . Atom $A2$ is prepared in state $|b_2\rangle$ and passes through cavity C . Atom $A3$ is prepared in state $|g_3\rangle$, and sent to the cavity C . Once the GHZ state has been obtained, the GHZ test is performed making use of the Ramsey cavities $K1, K2$ and $K3$ and detectors $D1, D2$ and $D3$ as described in the text.

References

- [1] M. Jammer, *The Philosophy of Quantum Mechanics. The Interpretation of Quantum Mechanics in Historical Perspective*, (John Wesley, New York, 1974).
- [2] R. Omnès, *The Interpretation of Quantum Mechanics*, (Princeton Univ. Press, Princeton, New Jersey, 1994).
- [3] R. I. G. Hughes, *The Structure and Interpretation of Quantum Mechanics*, (Harvard Univ. Press, Massachusetts, 1989).
- [4] H. Krips, *The Metaphysics of Quantum Theory*, (Oxford Univ. Press, Oxford, 1987).
- [5] B. d'Espagnat, *Scientific Am.* **241**, 128 (1979).
- [6] A. Einstein, B. Podolsky and N. Rosen, *Phys. Rev.* **47**, 777 (1935).
- [7] D. F. Walls and G. J. Milburn, *Quantum Optics*, (Springer-Verlag, Berlin, 1994).
- [8] N. Bohr, *Phys. Rev.* **48**, 696 (1935).
- [9] D. Bohm *Phys. Rev.* **85**, 166 (1952); D. Bohm *Phys. Rev.* **85**, 180 (1952); D. Bohm and J. Bud, *Rev. Mod. Phys.* **38**, 453 (1966).
- [10] J. S. Bell, *Physics* **1** (Long Island City, N. Y., 1964) 195; *Rev. Mod. Phys.* **38**, 447 (1966); *Speakable and Unsayable in Quantum Mechanics*, (Cambridge Univ. Press, Cambridge, 1987).
- [11] A. Aspect, P. Grangier and G. Roger, *Phys. Rev. Lett.* **47**, 460 (1981); *Phys. Rev. Lett.* **49**, 91 (1982); A. Aspect, J. Dalibard and G. Roger, *Phys. Rev. Lett.* **49**, 1804 (1982).
- [12] M. Hillery and B. Yurke, *Quantum Opt.* **7**, 215 (1995).
- [13] D. M. Greenberger, M. A. Horne and A. Zeilinger, in *Bell's Theorem, Quantum Theory, and Conceptions of the Universe*, Ed. M. Kafatos (Kluwer, Dordrecht, 1989) p. 74; D. M. Greenberger, M. A. Horne, A. Shimony and A. Zeilinger, *Am. J. Phys.* **58**, 1131 (1990).
- [14] N. D. Mermin, *Phys. Today* **43**, 9 (1990); *Am. J. Phys.* **58**, 731 (1990).
- [15] B. G. Englert and H. Walther, *Opt. Commun.* **179**, 283 (2000).
- [16] J. M. Raimond, M. Brune and S. Haroche, *Rev. Mod. Phys.* **73**, 565 (2001).
- [17] E. S. Guerra and J. C. Retamal, *J. Mod. Optics* **46**, 295 (1999).

- [18] C. C. Gerry, Phys. Rev. A **53**, 2857 (1996); Phys. Rev. A **54**, R2529 (1996).
- [19] D. Bouwmeester, J.-W. Pan, M. Daniell, H. Weinfurter and A. Zeilinger, Phys. Rev. Lett. **82**, 1345 (1999); J.-W. Pan, D. Bouwmeester, M. Daniell, H. Weinfurter and A. Zeilinger, Nature **403**, 515 (2000).
- [20] A. A. Radzig and B. M. Smirnov, *Reference Data on Atoms, Molecules, and Ions* (Springer-Verlag, Berlin, 1985); T. F. Gallagher, *Rydberg Atoms* (Cambridge Univ. Press, Cambridge, 1984).
- [21] M. Brune, F. Schmidt-Kaler, A. Maali, J. Dreyer, E. Hagley, J. M. Raimond, and S. Haroche, Phys. Rev. Lett. **76**, 1800 (1996).
- [22] G. Rempe, F. Schmidt-Kaler and H. Walther, Phys. Rev. Lett. **64**, 2783 (1990).
- [23] M. Orszag, *Quantum Optics*, (Springer-Verlag, Berlin, 2000).
- [24] P. L. Knight, Phys. Scr. **T12**, 51 (1986); S. J. D. Phoenix and P. L. Knight, J. Opt. Soc. Am. B **7**, 116 (1990).

Fig. 1 - E. S. Guerra




Fig. 2 - E. S. Guerra




Fig. 3 - E. S. Guerra




Fig. 4 - E. S. Guerra

